



POINTWISE CONVERGENCE FOR EXPANSIONS IN SPHERICAL MONOGENICS*

Fei Minggang (费铭岗)¹ Qian Tao (钱涛)²

1.School of Applied Mathematics, University of Electronic Science and Technology of China,
Changdu 610054, China

2.Department of Mathematics, Faculty of Science and Technology, University of Macau, Macao, China
E-mail: fei@uestc.edu.cn; fsttq@umac.mo

Abstract We offer a new approach to deal with the pointwise convergence of Fourier-Laplace series on the unit sphere of even-dimensional Euclidean spaces. By using spherical monogenics defined through the generalized Cauchy-Riemann operator, we obtain the spherical monogenic expansions of square integrable functions on the unit sphere. Based on the generalization of Fueter's theorem inducing monogenic functions from holomorphic functions in the complex plane and the classical Carleson's theorem, a pointwise convergence theorem on the new expansion is proved. The result is a generalization of Carleson's theorem to the higher dimensional Euclidean spaces. The approach is simpler than those by using special functions, which may have the advantage to induce the singular integral approach for pointwise convergence problems on the spheres.

Key words spherical monogenics; pointwise convergence of Fourier-Laplace series; generalized Cauchy-Riemann operator; unit sphere; generalization of Fueter's theorem

2000 MR Subject Classification 42B05; 30G35

1 Introduction

We start with reviewing the basic notations and results. Let $f \in L^1([-\pi, \pi])$, then the Fourier coefficients c_k are all well defined by

$$c_k = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(t) e^{-ikt} dt, \quad k \in \mathbf{Z}, \quad (1.1)$$

where \mathbf{Z} denotes the set of all integers. By $s_N(f)(x)$, we denote the partial sum

$$s_N(f)(x) = \sum_{|k| \leq N} c_k e^{ikx}, \quad x \in [-\pi, \pi], \quad N \in \mathbf{N}_0, \quad (1.2)$$

*Received August 23, 2007. Sponsored by Research Grant of the University of Macau No. RG024/03-04S/QT/FST

of the Fourier series of f , where \mathbf{N}_0 denotes the set of all natural numbers. Then we have,

$$s_N(f)(x) = \frac{1}{\pi} \int_{-\pi}^{\pi} f(t) D_N(x-t) dt, \quad (1.3)$$

where

$$D_N(x) = \begin{cases} \frac{\sin(N + \frac{1}{2})x}{2 \sin \frac{x}{2}} & \text{for } x \in [-\pi, \pi] \setminus \{0\}, \\ N + \frac{1}{2} & \text{for } x = 0 \end{cases}$$

is called the N -th Dirichlet kernel.

Since $L^2([-\pi, \pi]) \subset L^1([-\pi, \pi])$, the Fourier coefficients of L^2 -functions are well defined. There holds the following Carleson's theorem.

Theorem 1.1 [1] If $f \in L^2([-\pi, \pi])$, then

$$s_N(f)(x) \rightarrow f(x), \quad \text{a.e. } x \in [-\pi, \pi], \text{ as } N \rightarrow +\infty.$$

L. Carleson proved this theorem in 1966. The next year, R.A. Hunt [7] further extended this result to $f \in L^p([-\pi, \pi])$, $1 < p < \infty$.

One naturally asks what is the analogous result for the unit sphere S^n in the $(n+1)$ -dimensional Euclidean space \mathbf{R}_1^n , where

$$\mathbf{R}_1^n = \{x = x_0 + \underline{x} \mid x_0 \in \mathbf{R}, \underline{x} \in \mathbf{R}^n\}?$$

For any square integrable function f defined on S^n , denoted by $f \in L^2(S^n)$, there is an associated Fourier-Laplace series:

$$f \sim \sum_{k=0}^{\infty} f_k, \quad (1.4)$$

where f_k is the homogeneous spherical harmonics of degree k . There were comprehensive studies on convergence and summability of Fourier-Laplace series on the unit sphere of higher dimensional Euclidean spaces (see [8], [17], [20]). However, except for the two-dimensional case, pointwise convergence, being the initial motivation of harmonic analysis, may be said to be very little known. The case $n = 1$ seems to be the only well studied case ([1], [21]). Dirichlet ([2]) made the first detailed study for the case $n = 2$, on the so-called Laplace series. Koschmieder ([22]) studied the case $n = 3$. Roetman ([17]) considered the general cases, and, under certain conditions, reduced the convergence problem for $n = 2k + 1$ to $n = 1$; and $n = 2k + 2$ to $n = 2$. Among others, Meaney ([10]) and Yu ([23]) addressed some related topics, including the L^p cases.

The theory of Fourier-Laplace series is not facilitated with a complex structure like what is in the complex plane. All the known studies on Fourier-Laplace series are heavily dependent on the properties of spherical harmonics, especially Legendre polynomials. In this study, we provide a new approach based on Clifford algebras. Based on the induced complex structure we obtain integral expressions of partial sums in terms of Dirichlet kernels where no knowledge of special functions is involved. Our approach is based on generalizations of Fueter's theorem

(or inducing theorem, see below) on inducing monogenic and harmonic functions from those for one complex variable. In particular, the Dirichlet kernels (see Section 3) are induced from those for one complex variable by using inducing theorem. The convergence results obtained here are stronger than those obtained in [17] with simpler proofs, while [17] heavily relies on the special functions machinery. The applicability of the method adopted in this article is restricted to the even dimensional cases. That is because, the Fueter's theorem in those cases reduces to pointwise differentiation, and further reduces to the problems in complex plane. Our proofs are based on this pointwise differentiation approach. On the other hand, for the odd dimensional cases, Fueter's theorem reduces to computation on Fourier multiplier operators, with connections to complex plane; or alternatively to pointwise differentiation but based on the three-dimensional Euclidean space. The odd dimensional cases would require different ideas ([11], [13], [15]).

This article is a further development of [5] and [9]. In [5], the authors obtained the detailed expression of the Dirichlet kernels restricted on the unit sphere. Then, Riemann-Lebesgue theorem, localization principle, and a Dini's type pointwise convergence theorem are proved. In this article, based on the classical Carleson's theorem, we further obtain a weaker condition in terms of certain Sobolev spaces that guarantees the pointwise convergence of the Fourier-Laplace series (1.4) (see Section 4 for details). The result is a generalization of Carleson's theorem to the higher dimensional Euclidean spaces. We include here some propositions and results from [5] without proofs.

2 Preliminaries

We will be working with \mathbf{R}_1^n , the real-linear span of $\mathbf{e}_0, \mathbf{e}_1, \dots, \mathbf{e}_n$, where \mathbf{e}_0 is identical with 1 and $\mathbf{e}_i \mathbf{e}_j + \mathbf{e}_j \mathbf{e}_i = -2\delta_{ij}$. \mathbf{R}_1^n is embedded into the Clifford algebra $\mathbf{R}^{(n)}$ generated by e_1, \dots, e_n . A typical element in \mathbf{R}_1^n is denoted $x = x_0 + \underline{x}$, where $x_0 \in \mathbf{R}$ and $\underline{x} = x_1 \mathbf{e}_1 + \dots + x_n \mathbf{e}_n \in \mathbf{R}^n$. We will study \mathbf{R}_1^n -variable and Clifford-valued functions and the concepts of left- and right-monogenic functions are introduced via the generalized Cauchy-Riemann operator $D = \frac{\partial}{\partial x_0} \mathbf{e}_0 + \frac{\partial}{\partial x_1} \mathbf{e}_1 + \dots + \frac{\partial}{\partial x_n} \mathbf{e}_n$: a function f with continuous first order derivatives is said to be left-monogenic or right-monogenic if $Df = 0$ or $fD = 0$ in its domain, respectively. A function is said to be monogenic, if it is both left- and right-monogenic. The Cauchy kernel is denoted by $E(x) = \frac{1}{\omega_n} \frac{\bar{x}}{|x|^{n+1}}, x \in \mathbf{R}_1^n \setminus \{0\}$, where $\omega_n = \frac{2\pi^{\frac{n+1}{2}}}{\Gamma(\frac{n+1}{2})}$ is the surface area of the n -dimensional unit sphere S^n in \mathbf{R}_1^n . The Cauchy kernel is both left- and right-monogenic.

Let f^0 be a complex-valued function defined in an open set O in the upper-half complex plane. Write $f^0(z) = u(t, s) + iv(t, s)$, where u and v are real-valued.

Denote, for $x \in \vec{O}$,

$$\vec{f}^0(x) = u(x_0, |\underline{x}|) + \frac{\underline{x}}{|\underline{x}|} v(x_0, |\underline{x}|),$$

where $\vec{O} = \{x \in \mathbf{R}_1^n | (x_0, |\underline{x}|) \in O\}$.

\vec{f}^0 is said to be the induced function from f^0 , and \vec{O} is the induced set from O . The concept of intrinsic sets and functions naturally fit into our theory. In the complex plane \mathbf{C} , a set is said to be intrinsic if it is open and symmetric with respect to the real axis; and a

function f^0 an intrinsic function if it is defined in an intrinsic set and satisfies $\overline{f^0(z)} = f^0(\bar{z})$ within its domain (see [11]). In the notation $f^0 = u + iv$, the above condition is equivalent to require that u is even and v is odd in their second argument. In particular, $v(x_0, 0) = 0$, i.e., f^0 is real-valued if it is restricted to the real line in its domain.

Denote by τ the mapping

$$\tau(f^0) = \kappa^{-1} \Delta^{\frac{n-1}{2}} \vec{f}^0,$$

where $\Delta = D\bar{D}$, $\bar{D} = \frac{\partial}{\partial x_0} \mathbf{e}_0 - \frac{\partial}{\partial x_1} \mathbf{e}_1 - \cdots - \frac{\partial}{\partial x_n} \mathbf{e}_n$ and $\kappa_n = (2i)^{n-1} \Gamma^2(\frac{n+1}{2})$ are the normalizing constant that makes $\tau((\cdot)^{-1}) = E$ (see [14]).

The operator $\Delta^{\frac{n-1}{2}}$ is defined via the Fourier multiplier transformation on tempered distributions $M : \mathcal{S}' \rightarrow \mathcal{S}'$ induced by the multiplier $m(\xi) = (2\pi i |\xi|)^{n-1}$:

$$Mf = \mathcal{R}(m\mathcal{F}f),$$

where

$$\mathcal{F}f(\xi) = \int_{\mathbf{R}_1^n} e^{2\pi i \langle x, \xi \rangle} f(x) dx,$$

and

$$\mathcal{R}h(x) = \int_{\mathbf{R}_1^n} e^{-2\pi i \langle \xi, x \rangle} h(\xi) d\xi.$$

It is noted that both the Fourier transformation \mathcal{F} and its inverse \mathcal{R} are defined on tempered distributions via pairing with rapidly decreasing functions.

If n is an odd integer, then $\Delta^{\frac{n-1}{2}}$ reduces to a point-wise differential operator that was first studied by M. Sce who extended Fueter's result to \mathbf{R}_1^n for odd $n \in \mathbf{Z}^+$ ([18]). The corresponding result for n even was obtained and discussed by [13] and [14]. We have the following ([6], [18], [12–14]).

Proposition 2.1 Let $f^0(z) = u(s, t) + iv(s, t)$ be an intrinsic function defined on an intrinsic set $O \subset \mathbf{C}$. Then, the function $\tau(f^0)$ is monogenic in \vec{O} .

If we consider f^0 to be of the form z^k , $k \in \mathbf{Z}$, then we define the monomial functions by

$$P^{(-k)} = \tau((\cdot)^{-k}), \quad P^{(k-1)} = I(P^{(-k)}), \quad k \in \mathbf{Z}^+,$$

where \mathbf{Z}^+ is the set of positive integers and I is the Kelvin inversion defined by

$$I(f)(x) = E(x)f(x^{-1}).$$

When n is odd, the striking facts ([14]) are

$$\tau((\cdot)^{(k)}) = P^{(k)}, \quad k = -1, -2, \dots; \quad (2.1)$$

$$\tau((\cdot)^{(k+n-1)}) = P^{(k)}, \quad k = 0, 1, 2, \dots. \quad (2.2)$$

We note that, for any k ,

$$P^{(k)}(y^{-1}x)E(y)$$

is monogenic both in x and y ([14]). Since $E(y)n(y) = 1$ on the sphere, by Cauchy's theorem, we have

$$\int_{S^n} P^{(k)}(y^{-1}x) d\sigma(y) = 0, \quad |x| = 1, \quad k \neq 0; \quad (2.3)$$

and, since $P^{(0)} = I(P^{(-1)}) = I(E) = 1$, we have

$$\frac{1}{\omega_n} \int_{S^n} P^{(0)}(y^{-1}x) d\sigma(y) = 1, \quad |x| = 1. \quad (2.4)$$

3 Expansions of the Fourier-Laplace Series and Dirichlet Kernels

In the frame of Clifford algebras, we are enable to decompose spherical harmonics into sums of spherical monogenics ([4]). For any k -spherical harmonic f_k , as appeared in the Fourier-Laplace series of $f \in L^2(S^n)$ in (1.4), one has the unique decomposition

$$f_k = g_k + h_k, \quad (3.1)$$

where g_k is the restriction on the sphere of a left-monogenic function of homogeneity k , and h_k the same restriction of a left-monogenic function of homogeneity $-k + 1 - n$ ([3], [4]). Results in [14] imply that

$$g_k(x) = \frac{1}{\omega_n} \int_{S^n} P^{(k)}(y^{-1}x) f(y) d\sigma(y),$$

and

$$h_k(x) = \frac{1}{\omega_n} \int_{S^n} P^{(-k)}(y^{-1}x) f(y) d\sigma(y).$$

Note that the expressions $P^{(k)}(y^{-1}x)$ make sense as the domain of $P^{(k)}$ may be extended to products of vectors ([14]). The partial sum corresponding to (1.4), denoted by $S_N f(x)$, may be expressed as

$$S_N f(x) = \frac{1}{\omega_n} \int_{S^n} D_N^{(n+1)}(y^{-1}x) f(y) d\sigma(y), \quad (3.2)$$

where

$$D_N^{(n+1)}(x) = \sum_{|k| \leq N} P^{(k)}(x) \quad (3.3)$$

is called the N -th Dirichlet kernel in \mathbf{R}_1^n . When n is an odd number, invoking (2.1) and (2.2),

$$D_N^{(n+1)}(x) = \tau \left(\sum_{k=-N}^{N+n-1} (\cdot)^{(k)} \right) (x). \quad (3.4)$$

In this article, we only consider n being odd numbers and therefore $(n-1)/2$ being positive integers. The computation of $D_N^{(n+1)}(x)$ is based on the inducing theorem obtained in [15], as cited in the following proposition.

Proposition 3.1 Let $h(t, s)$ be harmonic for t and s in a region O in which $s > 0, t > 0$. Let $n \in \mathbf{Z}^+$ being odd and $x = x_0 + \underline{x} \in \mathbf{R}_1^n$. Define

$$H(x) = \Delta^{(n-1)/2} h(x_0, |\underline{x}|),$$

where the Laplacian Δ is in the $n+1$ variables, then

$$H(x) = (n-1)!! \left(\frac{1}{s} \partial_s \right)^{(n-1)/2} h(t, s) |_{t=x_0, s=|\underline{x}|}, \quad (3.5)$$

and H is harmonic in x_0, x_1, \dots, x_n in the corresponding region in \mathbf{R}_1^n .

Without loss of generality, we assume that functions to be expanded in (1.4) are scalar-valued. Indeed, a Clifford-valued function may be separated into 2^n parts of each which is scalar-valued.

Assume that $f \in L^2(S^n)$ and f is associated with an expansion (1.4). From the decomposition (3.1) and the integral formulas for g_k and h_k , we have

$$f_k(x) = \frac{1}{\omega_n} \int_{S^n} (P^{(k)} + P^{(-k)})(y^{-1}x)f(y)d\sigma(y).$$

Since f is scalar-valued, the scalar part of $P^{(k)} + P^{(-k)}$ will produce f_k , and the nonscalar parts of $P^{(k)}$ and $P^{(-k)}$ will have to be canceled out. This concludes that only the scalar parts of the Dirichlet kernels are concerned. We denote the scalar part of $D_N^{(n+1)}$ by $\mathcal{D}_N^{(n+1)}$.

Denote

$$f_N^0(z) = \sum_{k=-N}^{N+n-1} z^k = z^{-N} + \cdots + z^{-1} + 1 + z + \cdots + z^{N+n-1},$$

and $f_N^0(z) = U_N(t, s) + iV_N(t, s)$, where $z = t + is$, $t, s \in \mathbf{R}$. Thus, both U_N and V_N are harmonic functions in t and s .

Since we only consider n being odd numbers, we write $n = 2l + 1, l \in \mathbf{Z}^+$ for the rest of the article. Then $\mathcal{D}_N^{(n+1)} = \mathcal{D}_N^{(2l+2)}$. From [5], we have the detailed expression of $\mathcal{D}_N^{(2l+2)}$.

Proposition 3.2 For any $x \in S^n$, i.e., $x = x_0 + x_1 + \cdots + x_n$, and $|x| = 1$. Then

$$\begin{aligned} \mathcal{D}_N^{(2l+2)}(x) &= \kappa_n^{-1}(2l)!! \sum_{j=1}^l \sum_{i=1}^j C_l^j(\sin \theta)^{j-2l} Q_i^{(j)}(\sin \theta, \cos \theta) \left(\frac{\partial}{\partial \theta}\right)^i U_N \\ &\quad + \kappa_n^{-1}(2l)!! \sum_{j=1}^l \sum_{i=1}^j C_l^j(\sin \theta)^{j-2l} R_i^{(j)}(\sin \theta, \cos \theta) \left(\frac{\partial}{\partial \theta}\right)^i V_N, \end{aligned} \quad (3.6)$$

where $\theta = \arccos x_0$, $Q_i^{(j)}$ and $R_i^{(j)}$ are homogeneous polynomials of degree j in two variables, $C_l^j = (-1)^{l-j} \frac{(2l-j-1)!(2l-2j-1)!!}{(j-1)!(2l-2j)!}$, $1 \leq j \leq l-1$; $C_l^l = 1$.

Now, we work out U_N and V_N in polar coordinates. We have

$$\begin{aligned} f_N^0(z) &= \sum_{k=-N}^{N+n-1} z^k = \frac{1}{z^N(1-z)} - \frac{z^{N+n}}{1-z} \\ &= \frac{\cos N\theta - r \cos(N+1)\theta - r^{2N+n} \cos(N+n)\theta + r^{2N+n+1} \cos(N+n-1)\theta}{r^N(r^2 - 2r \cos \theta + 1)} \\ &\quad + i \frac{-\sin N\theta + r \sin(N+1)\theta - r^{2N+n} \sin(N+n)\theta + r^{2N+n+1} \sin(N+n-1)\theta}{r^N(r^2 - 2r \cos \theta + 1)} \\ &= U_N(r \cos \theta, r \sin \theta) + iV_N(r \cos \theta, r \sin \theta). \end{aligned}$$

Restricting to the unit sphere, we have

$$\begin{aligned} U_N(\cos \theta, \sin \theta) &= \frac{\cos N\theta - \cos(N+1)\theta - \cos(N+n)\theta + \cos(N+n-1)\theta}{2(1 - \cos \theta)} \\ &= \frac{\sin \frac{2N+1}{2}\theta + \sin \frac{2N+2n-1}{2}\theta}{2 \sin \frac{\theta}{2}} = \frac{\sin(N + \frac{n}{2})\theta \cos \frac{n-1}{2}\theta}{\sin \frac{\theta}{2}}, \end{aligned}$$

and

$$\begin{aligned} V_N(\cos \theta, \sin \theta) &= \frac{-\sin N\theta + \sin(N+1)\theta - \sin(N+n)\theta + \sin(N+n-1)\theta}{2(1 - \cos \theta)} \\ &= \frac{\cos \frac{2N+1}{2}\theta - \cos \frac{2N+2n-1}{2}\theta}{2 \sin \frac{\theta}{2}} = \frac{\sin(N + \frac{n}{2})\theta \sin \frac{n-1}{2}\theta}{\sin \frac{\theta}{2}}. \end{aligned}$$

4 Main Results

From the study in Section 3, we know that the Dirichlet kernels $\mathcal{D}_N^{(n+1)}(x)$ on S^n depend only on the angle θ , where $\theta = \arccos x_0$, $\theta \in [0, \pi]$, which means that $\mathcal{D}_N^{(n+1)}(y^{-1}x)$ on S^n depend only on $\text{Re}(y^{-1}x)$. In addition, when $x, y \in S^n$, we have that $\text{Re}(y^{-1}x) = \text{Re}(\bar{y}x) = \langle y, x \rangle$. Now, let x be a fixed point on S^n and write $y = x \cos \theta + \tilde{y} \sin \theta$, where \tilde{y} is orthogonal to x and $\theta = \arg(y, x)$. In this case, $y^{-1} = \bar{y} = \bar{x} \cos \theta + \bar{\tilde{y}} \sin \theta$ and $\text{Re}(y^{-1}x) = \langle y, x \rangle = \cos \theta$. We take average of a function $f \in L^2(S^n)$ over the $(n-1)$ -dimensional sphere whose points y satisfy $\arg(y, x) = \theta$. This average is denoted by

$$\Phi_x(f)(\theta) = \frac{1}{\omega_{n-1}} \int_{S^{n-1}} f(x \cos \theta + \tilde{y} \sin \theta) d\sigma_{n-1}(\tilde{y}),$$

where \tilde{y} is the spherical variable on S^{n-1} , $d\sigma_{n-1}(\tilde{y})$ is the normalized surface area measure on S^{n-1} . We call $\Phi_x(f)(\theta)$ the average of f about x in angle θ .

Note that in considering convergence problems it suffices to assume f to be scalar-valued. Without repeating, we will always assume this for the rest of the article. We briefly write $\Phi_x(f)(\theta) = \Phi_x(\theta)$. In addition, we use $\mathcal{W}_2^{l,1}([0, \pi])$ to denote the Sobolev space

$$\mathcal{W}_2^{l,1}([0, \pi]) = \left\{ g \in L^2([0, \pi]) \mid \left(\frac{\partial}{\partial \theta} \right)^k g \in L^1([0, \pi]), \ k = 1, 2, \dots, l \right\}.$$

Denote $\Phi_x(0) = \lim_{\theta \rightarrow 0} \Phi_x(\theta)$ if the limit exists. Then, the pointwise convergence theorem is as follows.

Theorem 4.1 Let x be a fixed point on S^n . For any $f \in L^2(S^n)$, we assume that $\Phi_x(0)$ exists. In addition, if $\Phi_x(\theta) \in \mathcal{W}_2^{l,1}([0, \pi])$, then $\lim_{N \rightarrow \infty} S_N(f)(x) = \Phi_x(0)$. If, in particular, f is continuous at x , then $\lim_{N \rightarrow \infty} S_N(f)(x) = f(x)$.

Proof It suffices to show $\lim_{N \rightarrow \infty} (S_N(f)(x) - \Phi_x(0)) = 0$. From the relations (2.3) and (2.4), we have

$$\frac{1}{\omega_n} \int_{S^n} D_N^{(n+1)}(y^{-1}x) d\sigma(y) = 1.$$

It is reduced to show

$$\lim_{N \rightarrow \infty} \frac{1}{\omega_n} \int_{S^n} D_N^{(n+1)}(y^{-1}x) (f(y) - \Phi_x(0)) d\sigma(y) = 0.$$

The last integral is scalar-valued and thus is induced from the scalar part of the N -th Dirichlet kernel $\mathcal{D}_N^{(n+1)} = \mathcal{D}_N^{(2l+2)}$, viz.

$$\frac{1}{\omega_n} \int_{S^n} \mathcal{D}_N^{(2l+2)}(y^{-1}x) (f(y) - \Phi_x(0)) d\sigma(y).$$

Substituting the expression of $\mathcal{D}_N^{(2l+2)}(y^{-1}x)$ and writing the integral into an iterated integral, the above becomes

$$\begin{aligned} & \frac{1}{\omega_n} \int_0^\pi (\sin \theta)^{(2l+2-2)} \left[\kappa_n^{-1}(2l)!! \sum_{j=1}^l \sum_{i=1}^j C_l^j(\sin \theta)^{j-2l} Q_i^{(j)}(\sin \theta, \cos \theta) \left(\frac{\partial}{\partial \theta} \right)^i U_N \right] \\ & \cdot (\Phi_x(\theta) - \Phi_x(0)) d\theta \end{aligned}$$

$$\begin{aligned}
& + \frac{1}{\omega_n} \int_0^\pi (\sin \theta)^{(2l+2-2)} \left[\kappa_n^{-1}(2l)!! \sum_{j=1}^l \sum_{i=1}^j C_l^j (\sin \theta)^{j-2l} R_i^{(j)}(\sin \theta, \cos \theta) \left(\frac{\partial}{\partial \theta} \right)^i V_N \right] \\
& \cdot (\Phi_x(\theta) - \Phi_x(0)) d\theta \\
& = \frac{1}{\omega_n} \kappa_n^{-1}(2l)!! \sum_{j=1}^l \sum_{i=1}^j C_l^j \int_0^\pi \left(\frac{\partial}{\partial \theta} \right)^i U_N [(\sin \theta)^j Q_i^{(j)}(\sin \theta, \cos \theta) (\Phi_x(\theta) - \Phi_x(0))] d\theta \\
& + \frac{1}{\omega_n} \kappa_n^{-1}(2l)!! \sum_{j=1}^l \sum_{i=1}^j C_l^j \int_0^\pi \left(\frac{\partial}{\partial \theta} \right)^i V_N [(\sin \theta)^j R_i^{(j)}(\sin \theta, \cos \theta) (\Phi_x(\theta) - \Phi_x(0))] d\theta.
\end{aligned}$$

Taking integration by parts repeatedly, we have

$$\begin{aligned}
& \lim_{N \rightarrow \infty} (S_N(f)(x) - \Phi_x(0)) \\
& = \lim_{N \rightarrow \infty} \frac{1}{\omega_n} \kappa_n^{-1}(2l)!! \sum_{j=1}^l \sum_{i=1}^j (-1)^i C_l^j \int_0^\pi U_N(\cos \theta, \sin \theta) \left(\frac{\partial}{\partial \theta} \right)^i \\
& \cdot \left[(\sin \theta)^j Q_i^{(j)}(\sin \theta, \cos \theta) (\Phi_x(\theta) - \Phi_x(0)) \right] d\theta \\
& + \lim_{N \rightarrow \infty} \frac{1}{\omega_n} \kappa_n^{-1}(2l)!! \sum_{j=1}^l \sum_{i=1}^j (-1)^i C_l^j \int_0^\pi V_N(\cos \theta, \sin \theta) \left(\frac{\partial}{\partial \theta} \right)^i \\
& \cdot \left[(\sin \theta)^j R_i^{(j)}(\sin \theta, \cos \theta) (\Phi_x(\theta) - \Phi_x(0)) \right] d\theta \\
& = \lim_{N \rightarrow \infty} I_1 + \lim_{N \rightarrow \infty} I_2.
\end{aligned}$$

We first consider the second part of the above expression. Since $\Phi_x(\theta) \in \mathcal{W}_2^{l,1}([0, \pi])$, $\Phi_x(\theta) - \Phi_x(0)$ belongs to $\mathcal{W}_2^{l,1}([0, \pi])$. As a consequence, $(\frac{\partial}{\partial \theta})^i [(\sin \theta)^j R_i^{(j)}(\sin \theta, \cos \theta) (\Phi_x(\theta) - \Phi_x(0))]$, $1 \leq i \leq j$, are all integrable. Replacing V_N by $\frac{\sin(N+\frac{n}{2})\theta \sin \frac{n-1}{2}\theta}{\sin \frac{\theta}{2}}$ and applying the classical Riemann–Lebesgue lemma, we have $\lim_{N \rightarrow \infty} I_2 = 0$.

As for I_1 , since U_N is equal to $\frac{\sin(N+\frac{n}{2})\theta \cos \frac{n-1}{2}\theta}{\sin \frac{\theta}{2}}$, there will be a singular point $\theta = 0$, thus the classical Riemann–Lebesgue lemma cannot be directly used. However, for any j , $1 \leq j \leq l$, we first consider the second summation $\sum_{i=1}^j$. There is a factor $(\sin \theta)^j$ in the integrand and the order of $\frac{\partial}{\partial \theta}$ is from 1 to j . When $1 \leq i \leq j-1$, after taking $(\frac{\partial}{\partial \theta})^i$ on $(\sin \theta)^j Q_i^{(j)}(\sin \theta, \cos \theta) (\Phi_x(\theta) - \Phi_x(0))$, there must be a factor $(\sin \theta)^{m_1}$ left, $m_1 \geq 1$. If $i = j$, after taking $(\frac{\partial}{\partial \theta})^j$ on $(\sin \theta)^j Q_i^{(j)}(\sin \theta, \cos \theta) (\Phi_x(\theta) - \Phi_x(0))$, there also must be a factor $(\sin \theta)^{m_2}$ left, $m_2 \geq 1$, except the first item that is $j!(\cos \theta)^j Q_i^{(j)}(\sin \theta, \cos \theta) (\Phi_x(\theta) - \Phi_x(0))$.

By using the classical Riemann–Lebesgue lemma for I_1 except the first items of the summation $\sum_{i=1}^j$ of I_1 , we have

$$\begin{aligned}
\lim_{N \rightarrow \infty} I_1 & = \lim_{N \rightarrow \infty} \frac{1}{\omega_n} \kappa_n^{-1}(2l)!! \sum_{j=1}^l (-1)^j C_l^j \int_0^\pi U_N(\cos \theta, \sin \theta) \\
& \cdot [j!(\cos \theta)^j Q_j^{(j)}(\sin \theta, \cos \theta) (\Phi_x(\theta) - \Phi_x(0))] d\theta \\
& = \lim_{N \rightarrow \infty} \frac{1}{\omega_n} \kappa_n^{-1}(2l)!! \sum_{j=1}^l (-1)^j j! C_l^j \int_0^\pi \frac{\sin(N+\frac{n}{2})\theta \cos \frac{n-1}{2}\theta}{\sin \frac{\theta}{2}}
\end{aligned}$$

$$\begin{aligned} & \cdot (\cos \theta)^j Q_j^{(j)}(\sin \theta, \cos \theta)(\Phi_x(\theta) - \Phi_x(0)) d\theta \\ &= \frac{2\pi}{\omega_n} \kappa_n^{-1} (2l)!! \sum_{j=1}^l (-1)^j j! C_l^j \lim_{N \rightarrow \infty} \frac{1}{\pi} \int_0^\pi \frac{\sin((N + \frac{n-1}{2}) + \frac{1}{2})\theta}{2 \sin \frac{\theta}{2}} \Psi_x^{(j)}(\theta) d\theta, \end{aligned}$$

where $\Psi_x^{(j)}(\theta) = \cos \frac{n-1}{2}\theta (\cos \theta)^j Q_j^{(j)}(\sin \theta, \cos \theta)(\Phi_x(\theta) - \Phi_x(0))$, $1 \leq j \leq l$. Since $\Phi_x(\theta) \in L^2([0, \pi])$, then, obviously, $\Psi_x^{(j)}(\theta) \in L^2([0, \pi])$, $1 \leq j \leq l$. By the classical Carleson's theorem, we obtain

$$\lim_{N \rightarrow \infty} I_1 = \frac{2\pi}{\omega_n} \kappa_n^{-1} (2l)!! \sum_{j=1}^l (-1)^j j! C_l^j \Psi_x^{(j)}(0) = 0.$$

Therefore, we have $\lim_{N \rightarrow \infty} S_N(f)(x) = \Phi_x(0)$.

Furthermore, by the result of R.A. Hunt ([7]), we can obviously extend the main convergence results to L^p cases, $1 < p < \infty$.

Now, we illustrate the results of Theorem 4.1 by two examples.

Example 4.1 It should be observed that for $n+1=2$, i.e., $l=0$ in Theorem 4.1, the average $\Phi_x(\theta)$ becomes the simple evaluation at two endpoints of the interval $(-\theta, \theta)$,

$$\Phi_x(\theta) = \frac{1}{2}[f(\theta_x + \theta) + f(\theta_x - \theta)],$$

where θ_x is the angle between x and e_1 , and the Dirichlet kernel $\mathcal{D}_N^{(2)}$ is just equal to D_N in Section 1. From the condition of Theorem 4.1, $\Phi_x \in L^2([0, \pi])$ if and only if $\frac{1}{2}[f(\theta_x + \theta) + f(\theta_x - \theta)] \in L^2([0, \pi])$. In particular, if $x=1$, Theorem 4.1 reduces to the classical Carleson's theorem.

Example 4.2 Let $l=1$, i.e., $n+1=4$. Theorem 4.1 becomes the following.

Let q be a fixed point on S^3 . For any $f \in L^2(S^3)$, we assume that $\Phi_q(0)$ exists. In addition, if $\Phi_q(\theta) \in L^2([0, \pi])$ and its derivative $\Psi_q(\theta)$ exists and $\Psi_q(\theta)$ is integrable in $(0, \pi)$, then $\lim_{N \rightarrow \infty} S_N(f)(q) = \Phi_q(0)$. Furthermore, $\lim_{N \rightarrow \infty} S_N(f)(q) = f(q)$ if f is continuous at q .

The convergence results in [9], with the assumption that $\Phi_q(\theta)$ is absolutely continuous in $[0, \pi]$, is a consequence of the above results. In fact, if $\Phi_q(\theta)$ is absolutely continuous in $[0, \pi]$, then $\Phi_q(\theta) \in L^2([0, \pi])$ and its derivative exists almost everywhere and is integrable on $[0, \pi]$.

For $l \geq 1$ the condition in Theorem 4.1 may be further weakened to the following:

$$\left\{ \begin{array}{l} \text{(i) } \cos \frac{n-1}{2}\theta (\cos \theta)^j Q_j^{(j)}(\sin \theta, \cos \theta)(\Phi_x(\theta) - \Phi_x(0)) \in L^2([0, \pi]), \quad 1 \leq j \leq l; \\ \text{(ii) } \left(\frac{\partial}{\partial \theta}\right)^i [(\sin \theta)^j Q_i^{(j)}(\sin \theta, \cos \theta)(\Phi_x(\theta) - \Phi_x(0))] \in L^1([0, \pi]), \quad 1 \leq i \leq j \leq l; \\ \text{(iii) } \left(\frac{\partial}{\partial \theta}\right)^i [(\sin \theta)^j R_i^{(j)}(\sin \theta, \cos \theta)(\Phi_x(\theta) - \Phi_x(0))] \in L^1([0, \pi]), \quad 1 \leq i \leq j \leq l. \end{array} \right.$$

Obviously, $\Phi_x(\theta) \in \mathcal{W}_2^{l,1}([0, \pi])$ implies the above conditions. The main results of [17] then can be easily deduced. In fact, there holds

Corollary 4.1 If f is differentiable up to order l on S^n , $l \geq 1$, i.e., $f \in C^l(S^n)$, then

$$\lim_{N \rightarrow \infty} S_N(f)(x) = f(x), \quad \forall x \in S^n.$$

Proof By Whitney's extension theorem ([19], also see [17]), we can show that $\Phi_x(\theta) \in C^l((0, \pi))$, and $(\sin \theta)^j \left(\frac{\partial}{\partial \theta}\right)^i \Phi_x(\theta)$, $1 \leq i \leq j \leq l$, converge to zero at 0 and π . In particular,

$\Phi_x(\theta)$ is continuous on $[0, \pi]$. Thus, we have that $\Phi_x(\theta) \in L^2([0, \pi])$ and $(\frac{\partial}{\partial \theta})^i[(\sin \theta)^j \Phi_x(\theta)]$ $1 \leq i \leq j \leq l$, are integrable in $[0, \pi]$. Then, the conditions (i), (ii), and (iii) are satisfied since $Q_i^{(j)}$ and $R_i^{(j)}$ are polynomials in $\sin \theta$ and $\cos \theta$. In addition, f is continuous on S^n , then we have,

$$\lim_{N \rightarrow \infty} S_N(f)(x) = f(x), \quad \forall x \in S^n.$$

References

- [1] Carleson L. On convergence and growth of partial sums of Fourier series. *Acta Math*, 1966, **116**: 135–157
- [2] Dirichlet P G L. Sur les séries dont le terme général dépend de deux angle, et qui servent á exprimer des fonctions arbitraires entre des limites données. *J Reine Angew Math*, 1873, **17**: 35–56
- [3] Delanghe R, Sommen F, Soucek V. Clifford algebra and spinor valued functions//A Function Theory for Dirac Operator. Dordrecht: Kluwer, 1992
- [4] Fei M G, Qian T. Direct sum decomposition of $L^2(\mathbf{R}_1^n)$ into subspaces invariant under Fourier transformation. *J Fourier Anal Appl*, 2006, **12**(2): 145–155
- [5] Fei M G, Qian T. Clifford algebra approach to pointwise convergence of Fourier series on spheres. *Science in China (Series A)*, 2006, **49**(11): 1553–1575
- [6] Fueter R. Die Funktionentheorie der Differentialgleichungen $\Delta u = 0$ und $\Delta \Delta u = 0$ mit vier reellen Variablen. *Comm Math Helv*, 1935, **7**: 307–330
- [7] Hunt R A. On the convergence of Fourier series, Orthogonal Expansions and Their continuous Analogues. *Proc Conf Edwardsville*, 111. 1967. 235–255; Southern Illinois Univ Press, Carbondale, 111. 1968
- [8] Kalf H. On the expansion of a function in terms of spherical harmonics in arbitray dimensions. *Bull Bel Math Soc*, 1995, **2**: 361–380
- [9] Liu S, Qian T. Pointwise convergence of Fourier series on the unit sphere of \mathbf{R}^4 with Quaternionic setting//Trends in Mathematics: Advance in Analysis and Geometry. Basel: Birkhäuser, 2004: 131–147
- [10] Meaney C. Divergence Jacobi polynomial series. *Proceedings of the American Mathematical Society*, 1983, **87**(3): 459–462
- [11] Peña D, Qian T, Sommen F. An alternative proof of Fueter’s theorem. *Complex Var Elliptic Equ*, 2006, **51**(8–11): 913–922
- [12] Qian T. Singular integrals on star-shaped Lipschitz surfaces in the quaternionic space. *Math Ann*, 1998, **310**(4): 601–630
- [13] Qian T. Generalization of Fueter’s result to \mathbf{R}^{n+1} . *Rend Mat Acc Lincei*, 1997, **8**(9): 111–117
- [14] Qian T. Fourier analysis on starshaped Lipschitz surfaces. *J of Func Anal*, 2001, **183**(2): 370–412
- [15] Qian T, Sommen F. Deriving harmonic functions in higher dimensional spaces. *Z Anal Anwendungen*, 2003, **22**(2): 275–288
- [16] Rinehart R F. Elements of theory of intrinsic functions on algebras. *Duke Math J*, 1965, **32**: 1–19
- [17] Roetman E L. Pointwise convergence for expansios in surface harmonics of arbitrary dimension. *J Reine Angew Math*, 1976, **282**: 1–10
- [18] Sce M. Osservazioni sulle serie di potenze nei moduli quadratici. *Atti Acc Lincei Rend fis*, 1957, **23**(8): 220–225
- [19] Whitney H. Differentiable functions defined in closed set I. *Trans Amer Math Soc*, 1934, **36**: 369–387
- [20] Wang K Y, Li L Q. Harmonic Analysis and Approximation on the Unit Sphere. Beijing, New York: Science Press, 2000
- [21] Zygmund A. Trigonometric Series, Vol 2. 2nd ed. Cambridge: Cambridge Univ Press, 1959
- [22] Koschmieder L. Unmittelbarer Beweis der Konvergenz einiger Riehen, die von mehreren Veränderlichen ab-hängen. *Monatsh Math*, 1934, **41**: 58–63
- [23] Yu J R. Some remarks on holomorphic functions and Taylor series in C^n . *Acta Math Sci*, 2008, **28B**(4): 721–726