

# Co-dimension- $p$ Shannon sampling theorems

YAN YANG\* and TAO QIAN

Faculty of Science and Technology, University of Macau, China

Communicated by F. Sommen

(Received 7 April 2006; in final form 26 April 2006)

In this article, by defining the generalized co-dimension- $p$  sinc function, the corresponding sinc interpolations (Shannon sampling theorems) are obtained.

*Keywords:* Shannon sampling; Sinc function; Clifford algebra

*AMS Subject Classifications:* 62D05, 30A05, 30D10, 42B35

## 1. Introduction

Sinc function in the real line  $\mathbf{R}$  is defined by

$$\text{sinc}(x) = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{ixt} dt = \frac{\sin(\pi x)}{\pi x}.$$

It has a holomorphic extension to the complex plane  $\mathbf{C}$ , i.e.,:

$$\text{sinc}(z) = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{izt} dt = \frac{\sin(\pi z)}{\pi z}. \quad (1)$$

Some applications of the sinc function may be found in [1].

For a set  $A$ , let  $\chi_A$  denote the characteristic function of  $A$ . Sinc function in  $\mathbf{R}^m$  is defined by

$$\begin{aligned} \text{sinc}(\underline{x}) &= (\chi_{[-\pi, \pi]^m})^\vee(\underline{x}) = \frac{1}{(2\pi)^m} \int_{\mathbf{R}^m} e^{i(\underline{x}, \underline{\xi})} \chi_{[-\pi, \pi]^m}(\underline{\xi}) d\underline{\xi} \\ &= \prod_{i=1}^m \text{sinc}(x_i) = \prod_{i=1}^m \frac{\sin(\pi x_i)}{\pi x_i}. \end{aligned}$$

---

\*Corresponding author. Email: ya27406@umac.mo

As a counterpart generalization of the sinc function (1) to higher-dimensional cases, Kou and Qian extended the sinc function in  $\mathbf{R}^m$  to  $m+1$ -dimensional real variables  $\mathbf{R}_1^m$  with the Clifford analysis setting in [2]. The definition of the extended sinc function (we call it *inhomogeneous co-dimension-1 sinc function*) is based on the generalized exponential function  $e(x, \underline{\xi})$  in  $\mathbf{R}_1^m \times \mathbf{R}^m$ , extending the classical exponential function  $e^{i(\underline{x}, \underline{\xi})}$  in  $\mathbf{R}^m \times \mathbf{R}^m$ . Furthermore, using the inhomogeneous co-dimension-1 sinc function, they obtained the Shannon sampling theorem [2] corresponding to the Paley–Wiener (P–W) theorem in  $\mathbf{R}_1^m$  [3]. In this article, we define co-dimension- $p$  sinc functions and prove the corresponding Shannon samplings in relation to the P–W theorems obtained in [4].

## 2. Preliminaries

For a basic knowledge and notation in relation to the Clifford algebra the readers are referred to [5–7].

Let  $\mathbf{e}_1, \dots, \mathbf{e}_m$  be *basic elements* satisfying  $\mathbf{e}_i \mathbf{e}_j + \mathbf{e}_j \mathbf{e}_i = -2\delta_{ij}$ , where  $\delta_{ij} = 1$  if  $i=j$  and  $\delta_{ij} = 0$  otherwise,  $i, j = 1, 2, \dots, m$ . Set

$$\mathbf{R}^m = \{\underline{x} = x_1 \mathbf{e}_1 + \dots + x_m \mathbf{e}_m : x_j \in \mathbf{R}, j = 1, 2, \dots, m\},$$

and

$$\mathbf{R}_1^m = \{x = x_0 + \underline{x} : x_0 \in \mathbf{R}, \underline{x} \in \mathbf{R}^m\}.$$

$\mathbf{R}^m$  and  $\mathbf{R}_1^m$  are called, respectively, the *homogeneous* and *inhomogeneous* Euclidean spaces.

Elements in  $\mathbf{R}^m$  are called *homogeneous vectors* and those of  $\mathbf{R}_1^m$  *inhomogeneous vectors* or *vectors*. The real (or complex) Clifford algebra generated by  $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_m$ , denoted by  $\mathbf{R}^{(m)}$  (or  $\mathbf{C}^{(m)}$ ), is the universal associative algebra generated by  $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_m$ , over the real (or complex) field  $\mathbf{R}$  (or  $\mathbf{C}$ ). A general element in  $\mathbf{R}^{(m)}$  (or  $\mathbf{C}^{(m)}$ ), a *Clifford number*, therefore, is of the form  $x = \sum_S x_S \mathbf{e}_S$ , where for  $S \neq \emptyset$ ,  $\mathbf{e}_S$  are *ordered reduced products* of the basis elements and  $\mathbf{e}_S = \mathbf{e}_{i_1} \mathbf{e}_{i_2} \dots \mathbf{e}_{i_l}$ , where  $S$  runs over all the ordered subsets of  $\{1, 2, \dots, m\}$ , namely

$$S = \{1 \leq i_1 < i_2 < \dots < i_l \leq m\}, \quad 1 \leq l \leq m,$$

and, for  $S = \emptyset$ , we set  $\mathbf{e}_\emptyset = \mathbf{e}_0 = 1$ .

The natural inner product between  $x$  and  $y$  in  $\mathbf{C}^{(m)}$ , denoted by  $\langle x, y \rangle$ , is the complex number  $\sum_S x_S \overline{y_S}$ , where  $x = \sum_S x_S \mathbf{e}_S$  and  $y = \sum_S y_S \mathbf{e}_S$ . The norm associated with this inner product is

$$|x| = \langle x, x \rangle^{1/2} = \left( \sum_S |x_S|^2 \right)^{1/2}.$$

The Clifford conjugate of a vector  $x = x_0 + \underline{x}$ , is defined to be  $\bar{x} = x_0 - \underline{x}$ . It is easy to verify that if  $x \neq 0$  and  $x \in \mathbf{R}_1^m$ , then  $x$  has an inverse,  $x^{-1}$ , and

$$x^{-1} = \frac{\bar{x}}{|x|^2}.$$

The *unit sphere*  $\{\underline{x} \in \mathbf{R}^m : |\underline{x}| = 1\}$  is denoted by  $S^{m-1}$ . We use  $B(\underline{x}, r)$  for the open ball in  $\mathbf{R}^m$  centered at  $\underline{x}$  with radius  $r$ , and  $\bar{B}(\underline{x}, r)$  for the topological closure of  $B(\underline{x}, r)$ .

Subsequently we shall study functions defined in the homogeneous space  $\mathbf{R}^m$  taking values in  $\mathbf{C}^{(m)}$ . So, they are of the form  $f(\underline{x}) = \sum_S f_S(\underline{x}) \mathbf{e}_S$ , where  $f_S$  are complex-valued functions. We shall use the *Dirac operator*, or the *homogeneous Dirac operator*,  $\partial_{\underline{x}}$ , where  $\partial_{\underline{x}} = (\partial/\partial x_1)\mathbf{e}_1 + \cdots + (\partial/\partial x_m)\mathbf{e}_m$ . We define the “left” and “right” roles of the operator  $\partial_{\underline{x}}$  by

$$\partial_{\underline{x}} f = \sum_{i=1}^m \sum_S \frac{\partial f_S}{\partial x_i} \mathbf{e}_i \mathbf{e}_S$$

and

$$f \partial_{\underline{x}} = \sum_{i=1}^m \sum_S \frac{\partial f_S}{\partial x_i} \mathbf{e}_S \mathbf{e}_i.$$

If  $\partial_{\underline{x}} f = 0$  in a domain (open and connected)  $U$ , then we say that  $f$  is *left-monogenic* in  $U$ ; and, if  $f \partial_{\underline{x}} = 0$  in  $U$ , then  $f$  is said to be *right-monogenic* in  $U$ . Left- or right-monogenic are called *one-sided-monogenic* or simply *monogenic*. The function theories for left- and right-monogenic functions, respectively, are parallel. If  $f$  is both left- and right-monogenic, then we say that  $f$  is *two-sided-monogenic*.

In  $\mathbf{R}_1^m$  we shall use the *inhomogeneous Dirac operator*, or the *generalized Cauchy–Riemann operator*,  $\partial_x = \partial_0 + \partial_{\underline{x}}$ ,  $\partial_0 = (\partial/\partial x_0)$ . The concept of monogenic functions in  $\mathbf{R}_1^m$  is defined via the inhomogeneous Dirac operator  $\partial_x$  in a similar manner. The monogenic function theories for the homogeneous and inhomogeneous spaces, respectively, are analogous.

Let  $k \in \mathbf{N}$ , where  $\mathbf{N}$  denotes the set of non-negative integers. Denote by  $M_\ell^+(m, k, \mathbf{C}^{(m)})$  the space of  $k$ -homogeneous left-monogenic polynomials in  $\mathbf{R}^m$ , whose restriction to  $S^{m-1}$  is denoted by  $\mathcal{M}_\ell^+(m, k, \mathbf{C}^{(m)})$ .

The Fourier transform of functions in  $\mathbf{R}^m$  is defined by

$$\hat{f}(\underline{\xi}) = \int_{\mathbf{R}^m} e^{-i(\underline{x}, \underline{\xi})} f(\underline{x}) d\underline{x},$$

and the inverse Fourier transform by

$$\check{g}(\underline{x}) = \frac{1}{(2\pi)^m} \int_{\mathbf{R}^m} e^{i(\underline{x}, \underline{\xi})} g(\underline{\xi}) d\underline{\xi},$$

where  $\underline{\xi} = \xi_1 \mathbf{e}_1 + \cdots + \xi_m \mathbf{e}_m$ .

To extend the domain of Fourier transforms to  $\mathbf{R}_1^m$ , we first need to extend the exponential function  $e^{i(\underline{x}, \underline{\xi})}$ . Denote, for  $x = x_0 \mathbf{e}_0 + \underline{x}$ ,

$$e(x, \underline{\xi}) = e^{i(\underline{x}, \underline{\xi})} e^{-x_0 |\underline{\xi}|} \chi_+(\underline{\xi}) + e^{i(\underline{x}, \underline{\xi})} e^{x_0 |\underline{\xi}|} \chi_-(\underline{\xi}), \quad (2)$$

where

$$\chi_{\pm}(\underline{\xi}) = \frac{1}{2} \left( 1 \pm i \frac{\underline{\xi} \mathbf{e}_0}{|\underline{\xi}|} \right).$$

It is easy to verify that the functions  $\chi_{\pm}$  satisfy the properties for projections:

$$\chi_- \chi_+ = \chi_+ \chi_- = 0, \quad \chi_{\pm}^2 = \chi_{\pm}, \quad \chi_+ + \chi_- = 1.$$

As extension of  $e(\underline{x}, \underline{\xi}) = e^{i(\underline{x}, \underline{\xi})}$  to  $\mathbf{R}_1^m \times \mathbf{R}^m$ , it is easy to verify that, for any fixed  $\underline{\xi}$ ,  $e(x, \underline{\xi})$  is two-sided-monogenic in  $x \in \mathbf{R}_1^m$ . The above extension is the inhomogeneous co-dimension-1 CK extension of  $e(\underline{x}, \underline{\xi})$  to  $\mathbf{R}_1^m$ . Replacing  $\mathbf{e}_0$  by  $\epsilon_0$  in equation (2), where  $\epsilon_0$  is a basis element added to the collection  $\mathbf{e}_1, \dots, \mathbf{e}_m$ , with  $\epsilon_0^2 = -1$  and anti-commutativity with the other  $\mathbf{e}_j, j = 1, \dots, m$ , one obtains the homogeneous co-dimension-1 CK extension  $e(x_0 \epsilon_0, \underline{x}, \underline{\xi})$  of  $e^{i(\underline{x}, \underline{\xi})}$  in  $\mathbf{R}^{m+1}$ . This function  $e(x_0 \epsilon_0, \underline{x}, \underline{\xi})$  is left-monogenic in  $x \in \mathbf{R}^{m+1}$ . Generalizations of the exponential function of these types can be first found in the work of Sommen [8], and then in Li *et al.* [7], where  $\underline{\xi}$  is further extended to  $\underline{\xi} + i\eta \in \mathbf{C}^m$ .

In [6], the generalized CK extension tells us: If  $A_0(\underline{y})$  is an analytic function in  $\mathbf{R}^q$ , for any  $k$ -homogeneous left-monogenic polynomial  $P_k(\underline{x})$  in  $\mathbf{R}^p$ , there exists a unique sequence  $(A_l(\underline{y}))_{l \geq 0}$  of analytic functions such that the series

$$\begin{aligned} f_{P_k}(\underline{x}, \underline{y}) &= \sum_{l=0}^{\infty} \underline{x}^l P_k(\underline{x}) A_l(\underline{y}) \\ &= \Gamma\left(k + \frac{p}{2}\right) \left( \frac{r \sqrt{\Delta_{\underline{y}}}}{2} \right)^{-(k+(p/2))} \left[ \frac{r \sqrt{\Delta_{\underline{y}}}}{2} J_{k+\frac{p}{2}-1}\left(r \sqrt{\Delta_{\underline{y}}}\right) + \frac{\underline{x} \partial_{\underline{y}}}{2} J_{k+(p/2)}\left(r \sqrt{\Delta_{\underline{y}}}\right) \right] (P_k(\underline{x}) A_0(\underline{y})), \end{aligned} \quad (3)$$

is convergent and its sum  $f_{P_k}$  is left-monogenic in any compact set belongs to  $\mathbf{R}^p \oplus \mathbf{R}^q$ . Where  $(\sqrt{\Delta_{\underline{y}}})^2 = \Delta_{\underline{y}}$ , the Laplacian in  $\underline{y}$ , and  $J_v$  the Bessel function

$$J_v(u) = \sum_{l=0}^{\infty} \frac{(-1)^l}{2^{2l+v} l! \Gamma(l+v+1)} u^{2l+v}.$$

We call  $f_{P_k}(\underline{x}, \underline{y})$  the *generalized CK extension in relation to  $P_k$  of  $A_0(\underline{y})$*  and  $A_0(\underline{y})$  the *initial value* of  $f_{P_k}(\underline{x}, \underline{y})$ . In particular, when  $k = 0, P_k = 1$ , we get  $A_0(\underline{y}) = f_{P_k}|_{\mathbf{R}^q}$ .

Denote  $\mathcal{T}_{P_k}(\mathbf{R}^q)$  the space of all functions of the form (3).

Furthermore, for any left-monogenic function  $f(\underline{x}, \underline{y})$  in  $\tilde{U}$  belongs to  $\mathbf{R}^m = \mathbf{R}^p \oplus \mathbf{R}^q$ , where  $\underline{x} = r\underline{\omega} \in \mathbf{R}^p$ ,  $\underline{y} \in \mathbf{R}^q$ , denote  $T_k(f)(\underline{\omega}, \underline{y}) = \lim_{r \rightarrow 0} 1/r^k P(k)f(r, \underline{\omega}, \underline{y})$ , where  $P(k)$  being the projection onto  $M_\ell^+(p, k, \mathbf{C}^{(p)})$ . It can be decomposed in variable  $\underline{x}$ . By using a basis  $P_{k,\alpha}(\underline{\omega})$  for  $M_\ell^+(p, k, \mathbf{C}^{(p)})$ ,

$$T_k(f)(\underline{\omega}, \underline{y}) = \sum_{\alpha \in A_k} P_{k,\alpha}(\underline{\omega}) T_{k,\alpha}(f)(\underline{y}),$$

where  $T_{k,\alpha}(f)(\underline{y})$  are real analytic functions.

Denote the generalized CK extension of  $T_{k,\alpha}(f)(\underline{y})$  with  $P_{k,\alpha}(\underline{x})$  by  $T_{k,\alpha}(\underline{x}, \underline{y})$ , then  $f$  can be written in a uniquely way as

$$f(\underline{x}, \underline{y}) = \sum_k \sum_{\alpha \in A_k} T_{k,\alpha}(\underline{x}, \underline{y}), \quad (4)$$

where  $T_{k,\alpha}(\underline{x}, \underline{y}) = \sum_l \underline{x}^l P_{k,\alpha}(\underline{x}) T_{k,\alpha}^{(l)}(f)(\underline{y})$  and the series (4) converging uniformly on any compact set in  $\tilde{U}$ . The series (4) is called *the generalized Taylor series* and  $T_{k,\alpha}^{(0)}(f)(\underline{y}) = T_{k,\alpha}(f)(\underline{y})$  are called *the initial values* of  $f$ .

To extend the domain of Fourier transforms to  $\mathbf{R}^p \oplus \mathbf{R}^q$ , we also need to extend the exponential function  $e^{i(\underline{y}, \underline{t})}$ . In [4], for a given  $k$ -homogeneous left-monogenic polynomial  $P_k(\underline{x})$ , we get the generalized CK extension of  $e^{i(\underline{y}, \underline{t})}$  in  $\mathcal{T}_{P_k}(\mathbf{R}^q)$ ,  $\underline{x} = r\underline{\omega} \in \mathbf{R}^p$ ,  $\underline{y}, \underline{t} \in \mathbf{R}^q$ , denoted by

$$\varepsilon_{P_k}^p(\underline{x}, \underline{y}, \underline{t}) = \Gamma\left(k + \frac{p}{2}\right) r^k e^{i(\underline{y}, \underline{t})} \left(\frac{r|\underline{t}|}{2}\right)^{-k-(p/2)+1} \left[ I_{k+(p/2)-1}(r|\underline{t}|) + i I_{k+(p/2)}(r|\underline{t}|) \underline{\omega} \frac{\underline{t}}{|\underline{t}|} \right] P_k(\underline{\omega}), \quad (5)$$

where

$$I_\nu(u) = i^{-\nu} J_\nu(iu) = \sum_{k=0}^{\infty} \frac{1}{k! \Gamma(\nu + k + 1)} \left(\frac{u}{2}\right)^{u+2k},$$

being a kind of Bessel functions. Then  $\varepsilon_{P_k}^p(\underline{x}, \underline{y}, \underline{t})$  is left-monogenic in  $\mathbf{R}^p \oplus \mathbf{R}^q$ .

In particular, when  $P_k = 1$ ,  $k = 0$ , we have

$$\varepsilon_1^p(\underline{x}, \underline{y}, \underline{t}) = \Gamma\left(\frac{p}{2}\right) e^{i(\underline{y}, \underline{t})} \left(\frac{r|\underline{t}|}{2}\right)^{-(p/2)+1} \left[ I_{(p/2)-1}(r|\underline{t}|) + i I_{p/2}(r|\underline{t}|) \underline{\omega} \frac{\underline{t}}{|\underline{t}|} \right].$$

We in particular denote

$$\varepsilon_1^1(x_1 \mathbf{e}_1, \underline{y}, \underline{t}) = e(x_1 \mathbf{e}_1, \underline{y}, \underline{t}).$$

From [4], we have, for any  $u > 0$ ,

$$\left(\frac{u}{2}\right)^{-\nu} I_\nu(u) = \sum_{k=0}^{\infty} \frac{1}{k! \Gamma(\nu + k + 1)} \left(\frac{u}{2}\right)^{2k} \leq C \sum_{k=0}^{\infty} \frac{u^{2k}}{(2k)!} \leq C e^u. \quad (6)$$

When  $|\underline{t}| \leq \Omega$ , we have

$$\begin{aligned} |\varepsilon_{P_k}^p(\underline{x}, \underline{y}, \underline{t})| &\leq C \left( \frac{2}{\Omega} \right)^k \left( \frac{r\Omega}{2} \right)^{-(p/2)+1} \left[ I_{k+\frac{p}{2}-1}(r\Omega) + I_{k+\frac{p}{2}}(r\Omega) \right] \\ &\leq C [I_k(r\Omega) + I_{k+1}(r\Omega)] \\ &\leq Ce^{r\Omega}. \end{aligned} \quad (7)$$

### 3. Exact interpolation with Shannon sampling in $\mathbf{R}^m = \mathbf{R}^p \oplus \mathbf{R}^q$

Based on the extension of the exponential function given by (5), the *generalized co-dimension- $p$  sinc function* in relation to  $P_k(\underline{x}) \in M_\ell^+(p, k, \mathbf{C}^{(p)})$  is defined by

$$\text{sinc}_{P_k}^p(\underline{x}, \underline{y}) = \frac{1}{(2\pi)^q} \int_{\mathbf{R}^q} \varepsilon_{P_k}^p(\underline{x}, \underline{y}, \underline{t}) \chi_{[-\pi, \pi]^q}(\underline{t}) d\underline{t}. \quad (8)$$

For  $h > 0$  fixed, define the *cardinal function* of  $f$  to be

$$C(f, h)(\underline{x}, \underline{y}) \equiv \sum_{\underline{k} \in \mathbf{Z}^q} \text{sinc}_{P_k}^p\left(\frac{\underline{x}}{h}, \frac{\underline{y} - h\underline{k}}{h}\right) f(h\underline{k}),$$

from equation (8), we have

$$\text{sinc}_{P_k}^p\left(\frac{\underline{x}}{h}, \frac{\underline{y} - h\underline{k}}{h}\right) = \frac{h^q}{(2\pi)^q} \int_{[-\pi/h, (\pi/h)^q]} \varepsilon_{P_k}^p(\underline{x}, \underline{y} - h\underline{k}, \underline{t}) d\underline{t}. \quad (9)$$

Next, we shall consider the generalized co-dimension- $p$  interpolation via the cardinal function corresponding to the generalized co-dimension- $p$  P-W theorem proved in [4]:

**LEMMA 1** [4] (Generalized co-dimension- $p$  P-W theorem) *Let  $P_k \in M_\ell^+(p, k; \mathbf{C}^{(p)})$  be given,  $F$  analytic, defined in  $\mathbf{R}^q$ , taking values in  $\mathbf{C}^{(q)}$ , which is the complex Clifford algebra generated by  $\mathbf{e}_{\mathbf{p}+1}, \dots, \mathbf{e}_{\mathbf{p}+q}$ , and  $F \in L^2(\mathbf{R}^q)$ ,  $\Omega$  be a positive real number. Then the following two assertions are equivalent:*

<sup>1</sup>  $F$  has a homogeneous co-dimensional- $p$  generalized CK extension to  $\mathbf{R}^{p+q}$ , denoted by  $f_{P_k}$ , and there exists a constant  $C$  such that

$$|f_{P_k}(\underline{x}, \underline{y})| \leq Ce^{\Omega|\underline{x}|}, \text{ for any } \underline{x} \in \mathbf{R}^p, \underline{y} \in \mathbf{R}^q.$$

<sup>2</sup>  $\text{supp}(\hat{F}) \subset \overline{B}(0, \Omega)$ .

Moreover, if one of the above conditions holds, then we have

$$f_{P_k}(\underline{x}, \underline{y}) = \frac{1}{(2\pi)^q} \int_{\mathbf{R}^q} \varepsilon_{P_k}^p(\underline{x}, \underline{y}, \underline{\xi}) \hat{F}(\underline{\xi}) d\underline{\xi}, \text{ for any } \underline{x} \in \mathbf{R}^p, \underline{y} \in \mathbf{R}^q.$$

A function  $f$  in  $\mathbf{R}^p \oplus \mathbf{R}^q$  is said to be of *exponential type  $\Omega$*  if

$$|f(\underline{x}, \underline{y})| \leq Ce^{\Omega|\underline{x}|}, \text{ for any } \underline{x} \in \mathbf{R}^p, \underline{y} \in \mathbf{R}^q$$

holds.

For any  $h > 0$ , denote

$$PW_{T_{P_k}(\mathbf{R}^q)}(\pi/h) = \{f | f \in \mathcal{T}_{P_k}(\mathbf{R}^q) \text{ and of exponential type } \pi/h, \text{ the initial value } F \in L^2(\mathbf{R}^q) \text{ and taking values in } \mathbf{C}^{(q)}\}.$$

Particularly, taking  $k = 0$ ,  $P_k = 1$  and  $p = 1$ , we have

$$PW_{\mathbf{R}^{q+1}}(\pi/h) = \{f | f \text{ is left-monogenic in } \mathbf{R}^{q+1} \text{ and of exponential type } \pi/h, f|_{\mathbf{R}^q} \in L^2(\mathbf{R}^q), \text{ and } f|_{\mathbf{R}^q} \text{ taking values in } \mathbf{C}^{(q)}\}.$$

The following theorems characterize the functions in the P–W class  $PW_{T_{P_k}(\mathbf{R}^q)}(\pi/h)$ .

**THEOREM 1** *If  $f \in PW_{T_{P_k}(\mathbf{R}^q)}(\pi/h)$ , then for any  $\underline{x} \in \mathbf{R}^p, \underline{y} \in \mathbf{R}^q$ , we have*

$1^0$

$$f(\underline{x}, \underline{y}) = \frac{1}{h^q} \int_{\mathbf{R}^q} \text{sinc}_{P_k}^p\left(\frac{\underline{x}}{h}, \frac{\underline{y} - \underline{\xi}}{h}\right) F(\underline{\xi}) d\underline{\xi}.$$

$2^0$

$$\frac{1}{(2\pi)^q} \int_{[-\pi/h, \pi/h]^q} |\hat{F}(\underline{y})|^2 d\underline{y} = \int_{\mathbf{R}^q} |F(\underline{t})|^2 d\underline{t} = \sum_{\underline{k} \in \mathbf{Z}^q} |F(h\underline{k})|^2, \quad (10)$$

where  $F$  is the initial value of  $f$ .

*Proof*  $1^0$ : Since  $f \in PW_{T_{P_k}(\mathbf{R}^q)}(\pi/h)$ , according to Lemma 1, we have

$$\begin{aligned} f(\underline{x}, \underline{y}) &= \frac{1}{(2\pi)^q} \int_{B(0, \pi/h)} \varepsilon_{P_k}^p(\underline{x}, \underline{y}, \underline{t}) \hat{F}(\underline{t}) d\underline{t} \\ &= \frac{1}{(2\pi)^q} \int_{[-\pi/h, \pi/h]^q} \varepsilon_{P_k}^p(\underline{x}, \underline{y}, \underline{t}) \hat{F}(\underline{t}) d\underline{t} \\ &= \frac{1}{(2\pi)^q} \int_{\mathbf{R}^q} \varepsilon_{P_k}^p(\underline{x}, \underline{y}, \underline{t}) \chi_{[-\pi/h, \pi/h]^q}(\underline{t}) \hat{F}(\underline{t}) d\underline{t}. \end{aligned}$$

By Parseval's theorem and (9), the above is equal to

$$\begin{aligned} &\frac{1}{(2\pi)^q} \int_{\mathbf{R}^q} \left[ \varepsilon_{P_k}^p(\underline{x}, \underline{y}, \underline{t}) \chi_{[-\pi/h, \pi/h]^q}(\underline{t}) \right] \widehat{(\underline{\xi}) F(\underline{\xi})} d\underline{\xi} \\ &= \frac{1}{h^q} \int_{\mathbf{R}^q} \left( \int_{[-\pi/h, \pi/h]^q} \varepsilon_{P_k}^p(\underline{x}, \underline{y} - \underline{\xi}, \underline{t}) d\underline{t} \right) F(\underline{\xi}) d\underline{\xi} \\ &= \frac{1}{h^q} \int_{\mathbf{R}^q} \text{sinc}_{P_k}^p\left(\frac{\underline{x}}{h}, \frac{\underline{y} - \underline{\xi}}{h}\right) F(\underline{\xi}) d\underline{\xi}. \end{aligned}$$

2<sup>0</sup>: From Lemma 1, we have

$$F(\underline{t}) = \frac{1}{(2\pi)^q} \int_{B(0, \pi/h)} e^{i(\underline{y}, \underline{t})} \hat{F}(\underline{y}) d\underline{y} = \frac{1}{(2\pi)^q} \int_{[-\pi/h, \pi/h]^q} e^{i(\underline{y}, \underline{t})} \hat{F}(\underline{y}) d\underline{y}.$$

Considering the Fourier expansion of  $\hat{F}$  in the cube  $[-\pi/h, \pi/h]^q$ , we have

$$h^q F(h\underline{k}) = \frac{1}{(2R)^q} \int_{[-R, R]^q} e^{i\pi(\underline{x}, \underline{k})/R} \hat{F}(\underline{y}) d\underline{y} = c_k,$$

where  $R = \frac{\pi}{h}$ , and  $c_k$  are the Fourier coefficients of  $\hat{F}$ . The Plancherel theorem of Fourier series is

$$\int_{[-R, R]^q} |\hat{F}(\underline{y})|^2 d\underline{y} = (2R)^q \sum_{\underline{k} \in \mathbf{Z}^q} |c_k|^2,$$

and the Plancherel theorem on  $L^2$ -functions in  $\mathbf{R}^q$  reads

$$\int_{\mathbf{R}^q} |\hat{F}(\underline{y})|^2 d\underline{y} = \int_{[-R, R]^q} |\hat{F}(\underline{y})|^2 d\underline{y} = (2\pi)^q \int_{\mathbf{R}^q} |F(\underline{t})|^2 d\underline{t}.$$

So we have

$$\frac{1}{(2\pi)^q} \int_{[-\pi/h, \pi/h]^q} |\hat{F}(\underline{y})|^2 d\underline{y} = \int_{\mathbf{R}^q} |F(\underline{t})|^2 d\underline{t} = \sum_{\underline{k} \in \mathbf{Z}^q} |F(h\underline{k})|^2. \quad \blacksquare$$

**COROLLARY 1** *If  $f \in PW_{T_1(\mathbf{R}^q)}(\pi/h)$ , then for any  $\underline{x} \in \mathbf{R}^p$ ,  $\underline{y} \in \mathbf{R}^q$ , we have*

$$f(\underline{x}, \underline{y}) = \frac{1}{h^q} \int_{\mathbf{R}^q} \text{sinc}_1^p\left(\frac{\underline{x}}{h}, \frac{\underline{y} - \underline{\xi}}{h}\right) f(\underline{\xi}) d\underline{\xi}.$$

$$\frac{1}{(2\pi)^q} \int_{[-\pi/h, \pi/h]^q} |\hat{f}(\underline{y})|^2 d\underline{y} = \int_{\mathbf{R}^q} |f(\underline{t})|^2 d\underline{t} = \sum_{\underline{k} \in \mathbf{Z}^q} |f(h\underline{k})|^2.$$

From (8) and the co-dimension- $p$  P-W theorem, we can obtain that  $\text{sinc}_{P_k}^p(\underline{x}/h, \underline{y}/h)$  belongs to  $PW_{T_{P_k}(\mathbf{R}^q)}(\sqrt{q}\pi/h)$ . Furthermore, we can construct functions in  $PW_{T_{P_k}(\mathbf{R}^q)}(\sqrt{q}\pi/h)$  using the following Theorem.

**THEOREM 2** *Let  $P_k \in M_\ell^+(p, k; \mathbf{C}^{(p)})$  be given,  $F \in L^2(\mathbf{R}^q)$  and take values in  $\mathbf{C}^{(q)}$ . Then  $f \in PW_{T_{P_k}(\mathbf{R}^q)}(\sqrt{q}\pi/h)$ , where*

$$f(\underline{x}, \underline{y}) = h^q \int_{\mathbf{R}^q} \text{sinc}_{P_k}^p\left(\frac{\underline{x}}{h}, \frac{\underline{y} - \underline{\xi}}{h}\right) F(\underline{\xi}) d\underline{\xi}. \quad (11)$$



*Proof* Applying the Parseval's theorem to the right-hand side of equation (11), owing to equation (9), we have

$$\begin{aligned}
 f(\underline{x}, \underline{y}) &= \frac{h^q}{(2\pi)^q} \int_{\mathbf{R}^q} \left[ \text{sinc}_{P_k}^p \left( \frac{\underline{x}}{h}, \frac{\underline{y} - \underline{\xi}}{h} \right) \right] (-\underline{t}) \hat{F}(\underline{t}) d\underline{t} \\
 &= \frac{h^q}{(2\pi)^q} \int_{\mathbf{R}^q} h^{-q} \varepsilon_{P_k}^p(\underline{x}, \underline{y}, \underline{t}) \chi_{[-\pi/h, \pi/h]^q}(\underline{t}) \hat{F}(\underline{t}) d\underline{t} \\
 &= \frac{1}{(2\pi)^q} \int_{\mathbf{R}^q} \varepsilon_{P_k}^p(\underline{x}, \underline{y}, \underline{t}) \chi_{[-\pi/h, \pi/h]^q}(\underline{t}) \hat{F}(\underline{t}) d\underline{t} \\
 &= \frac{1}{(2\pi)^q} \int_{[-\pi/h, \pi/h]^q} \varepsilon_{P_k}^p(\underline{x}, \underline{y}, \underline{t}) \hat{F}(\underline{t}) d\underline{t}.
 \end{aligned}$$

According to the evaluation (7) of  $\varepsilon_{P_k}^p$ , we have

$$|f(\underline{x}, \underline{y})| \leq C e^{\sqrt{q}\pi/h|\underline{x}|}, \text{ for any } \underline{x} \in \mathbf{R}^p, \underline{y} \in \mathbf{R}^q.$$

By Lemma 1, we get  $f(\underline{x}, \underline{y}) \in PW_{T_{P_k}(\mathbf{R}^q)}(\sqrt{q}\pi/h)$ . ■

Next, the exact  $\text{sinc}_{P_k}$  interpolation of functions in  $PW_{T_{P_k}(\mathbf{R}^q)}(\pi/h)$  is given.

**THEOREM 3** *If  $f \in PW_{T_{P_k}(\mathbf{R}^q)}(\pi/h)$ , then for any  $\underline{x} \in \mathbf{R}^p$ ,  $\underline{y} \in \mathbf{R}^q$ ,*

$$f(\underline{x}, \underline{y}) = C(f, h)(\underline{x}, \underline{y}) = \sum_{\underline{k} \in \mathbf{Z}^q} \text{sinc}_{P_k}^p \left( \frac{\underline{x}}{h}, \frac{\underline{y} - h\underline{k}}{h} \right) F(h\underline{k}), \quad (12)$$

where  $F$  is the initial value of  $f$  and the series on the right-hand side is absolutely and uniformly convergent for any  $\underline{y} \in \mathbf{R}^q$  and  $\underline{x}$  belongs to any bounded set in  $\mathbf{R}^p$ .

*Proof* Since  $f(\underline{x}, \underline{y}) \in PW_{T_{P_k}(\mathbf{R}^q)}(\pi/h)$ , Lemma 1 gives

$$f(\underline{x}, \underline{y}) = \frac{1}{(2\pi)^q} \int_{[-\pi/h, \pi/h]^q} \varepsilon_{P_k}^p(\underline{x}, \underline{y}, \underline{t}) \hat{F}(\underline{t}) d\underline{t}. \quad (13)$$

Expanding  $\varepsilon_{P_k}^p(\underline{x}, \underline{y}, \underline{t})$  on the cube  $[-\pi/h, \pi/h]^q$  into its multiple Fourier series in  $q$ -variables, we have

$$\varepsilon_{P_k}^p(\underline{x}, \underline{y}, \underline{t}) = \sum_{\underline{k} \in \mathbf{Z}^q} e^{i(h\underline{k}, \underline{t})} a_{\underline{k}}(\underline{x}, \underline{y}), \quad (14)$$

where

$$\begin{aligned}
 a_{\underline{k}}(\underline{x}, \underline{y}) &= \frac{h^q}{(2\pi)^q} \int_{[-\pi/h, \pi/h]^q} \varepsilon_{P_k}^p(\underline{x}, \underline{y}, \underline{t}) e^{-i(h\underline{k}, \underline{t})} d\underline{t} \\
 &= \frac{h^q}{(2\pi)^q} \int_{[-\pi/h, \pi/h]^q} \varepsilon_{P_k}^p(\underline{x}, \underline{y} - h\underline{k}, \underline{t}) d\underline{t} \\
 &= \text{sinc}_{P_k}^p \left( \frac{\underline{x}}{h}, \frac{\underline{y} - h\underline{k}}{h} \right)
 \end{aligned}$$

are the Fourier coefficients of  $\varepsilon_{P_k}^p(\underline{x}, \underline{y}, \underline{t})$ .

Substituting the series expansion (14) in the integral (13) and interchanging the order of the summation and the integration due to the  $L^2$ -convergence, we have

$$\begin{aligned} f(\underline{x}, \underline{y}) &= \frac{1}{(2\pi)^q} \int_{[-\pi/h, \pi/h]^q} \sum_{\underline{k} \in \mathbf{Z}^q} e^{i(h\underline{k}, \underline{t})} \text{sinc}_{P_k}^p\left(\frac{\underline{x}}{h}, \frac{\underline{y} - h\underline{k}}{h}\right) \hat{F}(\underline{t}) d\underline{t} \\ &= \sum_{\underline{k} \in \mathbf{Z}^q} \text{sinc}_{P_k}^p\left(\frac{\underline{x}}{h}, \frac{\underline{y} - h\underline{k}}{h}\right) \left( \frac{1}{(2\pi)^q} \int_{[-\pi/h, \pi/h]^q} e^{i(h\underline{k}, \underline{t})} \hat{F}(\underline{t}) d\underline{t} \right) \\ &= \sum_{\underline{k} \in \mathbf{Z}^q} \text{sinc}_{P_k}^p\left(\frac{\underline{x}}{h}, \frac{\underline{y} - h\underline{k}}{h}\right) F(h\underline{k}). \end{aligned}$$

We next show the uniform convergence of the series on the right side.

In fact, for any positive number  $M$ , using the Cauchy–Schwarz inequality, we have

$$\left| \sum_{|\underline{k}| \leq M} \text{sinc}_{P_k}^p\left(\frac{\underline{x}}{h}, \frac{\underline{y} - h\underline{k}}{h}\right) F(h\underline{k}) \right| \leq \left( \sum_{|\underline{k}| \leq M} \left| \text{sinc}_{P_k}^p\left(\frac{\underline{x}}{h}, \frac{\underline{y} - h\underline{k}}{h}\right) \right|^2 \right)^{1/2} \left( \sum_{|\underline{k}| \leq M} |F(h\underline{k})|^2 \right)^{1/2}.$$

Note that the function  $\varepsilon_{P_k}^p(\underline{x}, \underline{y}, \cdot) \in L^2([-\pi/h, \pi/h]^q)$ . Using the Bessel inequality and equation (7), for any bounded set  $U \in \mathbf{R}^p$ , we have

$$\begin{aligned} \left( \sum_{|\underline{k}| \leq M} \left| \text{sinc}_{P_k}^p\left(\frac{\underline{x}}{h}, \frac{\underline{y} - h\underline{k}}{h}\right) \right|^2 \right)^{1/2} &\leq \left( \frac{h}{2\pi} \right)^{q/2} \|\varepsilon_{P_k}^p(\underline{x}, \underline{y}, \cdot)\|_{L^2([-\pi/h, \pi/h]^q)} \\ &\leq \left( \frac{h}{2\pi} \right)^{q/2} e^{\sqrt{q}|\underline{x}|\pi/h} \leq C < \infty, \end{aligned}$$

where  $\underline{y} \in \mathbf{R}^q$ ,  $\underline{x} \in U$ . Owing to the estimate and equation (10), the series in equation (12) is convergent uniformly and absolutely in  $U \oplus \mathbf{R}^q$ .  $\blacksquare$

The homogeneous co-dimension- $p$  P–W theorem is stated as:

**LEMMA 2** [4] (Homogeneous co-dimension- $p$  P–W theorem) *Let  $F$  be analytic, defined in  $\mathbf{R}^q$ , taking values in  $\mathbf{C}^{(q)}$ , the complex Clifford algebra generated by  $\mathbf{e}_{p+1}, \dots, \mathbf{e}_{p+q}$ , and  $F \in L^2(\mathbf{R}^q)$ .  $\Omega$  is a positive real number. Then the following two assertions are equivalent:*

$1^0$   *$F$  has a homogeneous co-dimensional- $p$  CK extension to  $\mathbf{R}^{p+q}$ , denoted by  $f$ , and there exists a constant  $C$  such that*

$$|f(\underline{x}, \underline{y})| \leq C e^{\Omega|\underline{x}|}, \text{ for any } \underline{x} \in \mathbf{R}^p, \underline{y} \in \mathbf{R}^q.$$

$2^0$   *$\text{supp}(\hat{F}) \subset \overline{B}(0, \Omega)$ .*

Moreover, if one of the above conditions holds, we have

$$f(\underline{x}, \underline{y}) = \frac{1}{(2\pi)^q} \int_{\mathbf{R}^q} \varepsilon_1^p(\underline{x}, \underline{y}, \underline{\xi}) \hat{F}(\underline{\xi}) d\underline{\xi}, \text{ for any } \underline{x} \in \mathbf{R}^p, \underline{y} \in \mathbf{R}^q.$$

Corresponding to the Lemma 2, we have the exact  $\text{sinc}_1^p$  interpolation of functions in  $PW_{T_1(\mathbf{R}^q)}(\pi/h)$ .

Corollary 2 *If  $f \in PW_{T_1(\mathbf{R}^q)}(\pi/h)$ , then for any  $\underline{x} \in \mathbf{R}^p, \underline{y} \in \mathbf{R}^q$ ,*

$$f(\underline{x}, \underline{y}) = C(f, h)(\underline{x}, \underline{y}) = \sum_{\underline{k} \in \mathbf{Z}^q} \text{sinc}_1^p\left(\frac{\underline{x}}{h}, \frac{\underline{y} - h\underline{k}}{h}\right) f(h\underline{k}),$$

where

$$\text{sinc}_1^p(\underline{x}, \underline{y}) = \frac{1}{(2\pi)^q} \int_{\mathbf{R}^q} \varepsilon_1^p(\underline{x}, \underline{y}, \underline{t}) \chi_{[-\pi, \pi]^q}(\underline{t}) d\underline{t},$$

and the series on the right-hand side is absolutely and uniformly convergent for any  $\underline{y} \in \mathbf{R}^q$  and  $\underline{x}$  belongs to any bounded set in  $\mathbf{R}^p$ .

Henceforth the article shall deal with the Shannon sampling theorem in relation to the generalized Taylor series.

In [4], the co-dimension-p P–W theorem related to generalized Taylor series reads:

LEMMA 3 *Assume that  $f(\underline{x}, \underline{y})$  is left-monogenic in  $\mathbf{R}^m = \mathbf{R}^p \oplus \mathbf{R}^q$  with the form (4). For any  $k \geq 0$  and  $\alpha \in A_k$ , let  $T_{k,\alpha}(f)(\underline{y}) = T_{k,\alpha}^{(0)}(f)(\underline{y})$  be analytic, defined in  $\mathbf{R}^q$ , taking values in  $\mathbf{C}^{(q)}$ , the complex Clifford algebra generated by  $\mathbf{e}_{p+1}, \dots, \mathbf{e}_{p+q}$ ,  $T_{k,\alpha}(f)(\underline{y}) \in L^2(\mathbf{R}^q)$ ,*

$$\left| \sum_k \sum_\alpha P_{k,\alpha}(\underline{x}) \hat{T}_{k,\alpha}(f)(\underline{\xi}) \right| \leq C e^{\Omega|\underline{x}|}, \text{ for any } \underline{x} \in \mathbf{R}^p, \underline{\xi} \in \mathbf{R}^q,$$

where  $\Omega$  is a positive real number. Then the following two assertions are equivalent:

1<sup>0</sup> *There exists a constant  $C$  such that*

$$|f(\underline{x}, \underline{y})| \leq C e^{\Omega|\underline{x}|}, \text{ for any } \underline{x} \in \mathbf{R}^p, \underline{y} \in \mathbf{R}^q.$$

2<sup>0</sup>  $\text{supp}(\hat{T}_{k,\alpha}(f)) \subset \overline{B}(0, \Omega)$ , for any  $k \geq 0$  and  $\alpha \in A_k$ .

Moreover, if one of the above conditions holds, we have

$$f(\underline{x}, \underline{y}) = \sum_k \sum_\alpha T_{k,\alpha}(\underline{x}, \underline{y}) = \frac{1}{(2\pi)^q} \sum_k \sum_\alpha \int_{\mathbf{R}^q} \varepsilon_{P_{k,\alpha}}^p(\underline{x}, \underline{y}, \underline{\xi}) \hat{T}_{k,\alpha}(f)(\underline{\xi}) d\underline{\xi}, \quad (15)$$

for any  $\underline{x} \in \mathbf{R}^p, \underline{y} \in \mathbf{R}^q$  and the series is converging uniformly on any compact set in  $\mathbf{R}^p \oplus \mathbf{R}^q$ .

Next, the Shannon sampling theorem corresponding to the P–W theorem above is obtained. For any  $h > 0$ , denote

$PW_{\mathbf{R}^p \oplus \mathbf{R}^q}(\pi/h) = \{f \mid f \text{ is left-monogenic in } \mathbf{R}^p \oplus \mathbf{R}^q \text{ with the form (4) and of exponential type } \pi/h, \text{ the initial values } T_{k,\alpha}(f)(\underline{y}) \in L^2(\mathbf{R}^q) \text{ and taking values in } \mathbf{C}^{(q)}.\}$

**THEOREM 4** *If  $f \in PW_{\mathbf{R}^p \oplus \mathbf{R}^q}(\pi/h)$ , then for any  $\underline{x} \in \mathbf{R}^p, \underline{y} \in \mathbf{R}^q$ ,*

$$f(\underline{x}, \underline{y}) = \sum_k \sum_\alpha C[T_{k,\alpha}(f), h](\underline{x}, \underline{y}), \quad (16)$$

where

$$C[T_{k,\alpha}(f), h](\underline{x}, \underline{y}) = \sum_{\underline{k} \in \mathbf{Z}^q} \text{sinc}_{P_{k,\alpha}}^p\left(\frac{\underline{x}}{h}, \frac{\underline{y} - h\underline{k}}{h}\right) T_{k,\alpha}(f)(h\underline{k}) \quad (17)$$

and  $T_{k,\alpha}(f)(\underline{y})$  are the initial values of  $f$ . The series (16) and (17) on the right-hand side are uniformly convergent on any compact set in  $\mathbf{R}^p \oplus \mathbf{R}^q$ .

*Proof* If  $f \in PW_{\mathbf{R}^p \oplus \mathbf{R}^q}(\pi/h)$ , then  $f$  has the form in (15). Using Theorem 3, we obtain

$$T_{k,\alpha}(\underline{x}, \underline{y}) = C[T_{k,\alpha}(f), h](\underline{x}, \underline{y}) = \sum_{\underline{k} \in \mathbf{Z}^q} \text{sinc}_{P_{k,\alpha}}^p\left(\frac{\underline{x}}{h}, \frac{\underline{y} - h\underline{k}}{h}\right) T_{k,\alpha}(f)(h\underline{k}). \quad \blacksquare$$

## Acknowledgement

This work was supported by Research Grant of the University of Macau No. RG024/03-04S/QT/FST.

## References

- [1] Lund, J. and Bowers, K., 1992, *Sinc Methods for Quadrature and Differential Equations*, SIAM.
- [2] Kou, K. and Qian, T., 2005, Shannon sampling with the Clifford Analysis setting. *Zeitschrift für Analysis und ihre Anwendungen*, **24**(4), 853–870.
- [3] Kou, K. and Qian, T., 2002, The Paley–Wiener theorem in  $\mathbf{R}_1^n$  with the Clifford analysis setting. *Journal of Functional Analysis*, **189**, 227–241.
- [4] Yang, Y., Qian, T. and Sommen, F., *Co-dimension-P Paley-Wiener Theorems* (preprint: Singapore).
- [5] Delanghe, R., Sommen, F. and Souček, V., 1992, *Clifford Algebra and Spinor-Valued Functions*, Vol. 53, (Dordrecht, Boston, London: Kluwer Academic Publishers).
- [6] Delanghe, R., Sommen, F. and Souček, V., 1992, Residues in Clifford analysis. *Partial Differential Equations with Complex Analysis*, Pitman Research Notes in Math., Vol. 262, pp. 61–92.
- [7] Li, C., McIntosh, A. and Qian, T., 1994, Clifford algebras, fourier transforms, and singular convolution operators on lipschitz surfaces. *Rev. Mat. Iberoamericana*, **10**, 665–721.
- [8] Sommen, F., 1982, Some connections between Clifford analysis and complex analysis. *Complex Variables: Theory Appl.*, **1**, 97–118.

Copyright of Complex Variables & Elliptic Equations is the property of Taylor & Francis Ltd and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.